

Can spectral and local information decide separability?

K. G. H. Vollbrecht* and M. M. Wolf†

Institut für Mathematische Physik, TU Braunschweig,
Mendelssohnstr.3, 38106 Braunschweig, Germany.
(February 1, 2008)

We discuss the discriminating power of separability criteria, which are based on the spectrum of a quantum state and its reductions. Common examples are entropic inequalities utilizing conditional Tsallis or Rényi entropies. We prove that these inequalities are implied by the reduction criterion for any positive value of the entropic parameters. We show however, that arbitrary sets of criteria based on spectral and local information can never be sufficient by establishing a separable, isospectral and locally undistinguishable counterpart for any Werner state in odd dimensions. For the case of two qubit systems we show that a simple controlled phase gate operation can produce an isospectral, entangled state out of a separable one, which has the same reductions.

03.65.Bz, 03.67.-a

I. INTRODUCTION

Entanglement has always been a key issue in the ongoing debate about the foundations and interpretation of quantum mechanics since Einstein [1] and Schrödinger [2] expressed their deep dissatisfaction about this astonishing part of quantum theory. Whereas for the long period from 1935 to 1964, until Bell [3] published his famous work, discussions about entanglement were purely meta-theoretical, nowadays quantum information theory has established entanglement as a physical resource and key ingredient for quantum computation and quantum information processing. This new point of view led to a dramatic increase of general structural knowledge about entanglement in the last few years. However, there are still many open problems and deciding them often fails at what we try to profit from in quantum computation: the rapid increase of dimensions. One of these unsolved problems is in fact the question whether a given quantum state is entangled or merely classically correlated, i.e. separable.

The present paper is devoted to the question to what extent the combination of spectral and local information can decide separability. We will show that the discriminating power of any separability criterion based on this

kind of information is rather limited or even null for some otherwise simple classes of quantum states.

The paper is organized as follows. In Sec. II we will discuss entropic inequalities as common examples for the kind of criteria we want to address. We give a small counterexample for the conjectured monotonicity of the conditional Tsallis entropy and prove that the criteria based on conditional Tsallis and Rényi entropies are implied by the reduction criterion for any positive value of the entropic parameters. In Sec. III, IV we will construct pairs of *isospectral* states having the same spectra and the same reductions such that one state is entangled and the other is separable. Sec. III establishes such an isospectral and separable counterpart for any entangled Werner state in odd dimensions. Sec. IV then provides a similar example even for the case of two qubits by utilizing a phase gate operation showing clearly how weak the discriminating power of the considered class of separability criteria is.

To fix ideas we will start by recalling some of the basic notions and previous results.

A bipartite quantum state described by its density matrix ρ acting on a Hilbert space $\mathcal{H} = \mathcal{H}^{(1)} \otimes \mathcal{H}^{(2)}$ is said to be *separable*, unentangled or classically correlated if it can be written as a convex combination of tensor product states [4]

$$\rho = \sum_j p_j \rho_j^{(1)} \otimes \rho_j^{(2)}, \quad (1)$$

where the positive weights p_j sum up to one and $\rho^{(i)}$ describes a state on $\mathcal{H}^{(i)}$. This means in particular that pure states are separable if and only if they are product states. Moreover, all entanglement properties of pure states, which can always be written in their Schmidt form (cf. [5]) as $|\Psi\rangle = \sum_i \sqrt{\lambda_i} |i\rangle \otimes |i\rangle$, are completely determined by the eigenvalues $\{\lambda_i\}$ of the reduced state $\rho_A = \text{tr}_B |\Psi\rangle\langle\Psi|$.

For mixed quantum states however, the situation is much more difficult and deciding whether a state is entangled or separable is not yet feasible in general. Currently, the most efficient necessary criterion for separability is the positivity of the partial transpose (PPT), i.e., the condition that ρ^{T_1} has to be a positive semi-definite operator [6]. The partial transpose of the state is thereby defined in terms of its matrix elements with respect to some basis by $\langle kl | \rho^{T_1} | mn \rangle = \langle ml | \rho | kn \rangle$. For the smallest non-trivial systems with 2×2 resp. 2×3 dimensional Hilbert spaces and a few other special cases the PPT-criterion also turned out to be sufficient [7]. In higher

*Electronic Mail: k.vollbrecht@tu-bs.de

†Electronic Mail: mm.wolf@tu-bs.de

dimensional systems however, so called *bound entangled* states exist, which satisfy the PPT-condition without being separable [8].

Another well known condition is given by the *reduction criterion* [9,10]

$$\rho_A \otimes \mathbf{1} - \rho \geq 0, \quad (2)$$

which is implied by the PPT-criterion but nevertheless an important condition since its violation implies the possibility of recovering entanglement by distillation (which is yet unclear for PPT violating states). For the case of two qubits the reduction criterion is also known to be sufficient [9]. The general line of implication is

$$\rho \text{ sep.} \Rightarrow \rho^{T_1} \geq 0 \Rightarrow \rho \text{ undistillable} \Rightarrow \rho_A \otimes \mathbf{1} - \rho \geq 0.$$

II. ENTROPIC INEQUALITIES

The idea to use entropic inequalities as separability criteria for mixed states goes back to the mid nineties when Cerf and Adami [11] and the Horodecki family [12] recognized that certain conditional Rényi entropies are non-negative for separable states, and it was recently resurrected by several groups [13–17] in the form of conditional Tsallis entropies. Characteristic for these criteria is, that they just use the spectra of the state and its reductions.

The quantum Rényi entropy depending on the parameter $\alpha \in \mathbb{R}$ is given by

$$S_\alpha(\rho) = \frac{\log \text{tr}(\rho^\alpha)}{1 - \alpha}. \quad (3)$$

For the case of separable states it was shown in [11,12,18]:

$$S_\alpha(\rho) - S_\alpha(\rho_A) \geq 0 \quad (4)$$

for $\alpha = 0, \infty$ and $\alpha \in [1, 2]$, where S_0, S_1, S_∞ reduces to the logarithm of the rank, the von Neumann entropy and the negative logarithm of the operator norm respectively.

In Ref. [13,15] essentially the same criterion was expressed in terms of the conditional Tsallis entropy

$$T_\alpha(\rho) = \frac{\text{tr}(\rho_A^\alpha) - \text{tr}(\rho^\alpha)}{(\alpha - 1) \text{tr}(\rho_A^\alpha)}, \quad (5)$$

which was conjectured [15] to be monotone in α and non-negative for all positive values of α if and only if ρ is separable.

However, it is definitively neither monotone nor sufficient for separability. Monotonicity can most easily be ruled out by low rank examples like

$$\rho = \frac{1}{2}(|\Phi_+\rangle\langle\Phi_+| + |01\rangle\langle 01|), \quad |\Phi_+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle),$$

for which the reduced state has eigenvalues $\frac{1}{4}, \frac{3}{4}$ and therefore $T_0 = T_\infty = 0 \neq T_2 = \frac{1}{5}$. Fortunately however, monotonicity is not necessary for proving the validity of the Tsallis/Rényi separability criteria for other values than $\alpha = 0, \infty, \alpha \in [1, 2]$ [19]:

Proposition 1 *Any state for which the reduction criterion (2) holds, in particular any separable state, satisfies*

$$\text{tr}(\rho_A^\alpha) - \text{tr}(\rho^\alpha) \begin{cases} \geq 0, & \alpha > 1 \\ \leq 0, & 0 \leq \alpha < 1 \end{cases}. \quad (6)$$

This implies the validity of the Tsallis/Rényi separability criteria for any positive value of the entropic parameter.

Proof : For $\alpha > 1$ the proof is essentially based on the Golden-Thompson inequality (cf. [20]) stating that $\text{tr}(e^A e^B) \geq \text{tr}(e^{A+B})$ for hermitian matrices A, B . Utilizing the definition of the reduced state, i.e., $\forall P \geq 0$: $\text{tr}(\rho(P \otimes \mathbf{1})) \equiv \text{tr}(\rho_A P)$ this leads to:

$$\text{tr}(\rho_A^\alpha) = \text{tr}[\rho(\rho_A^{\alpha-1} \otimes \mathbf{1})] \quad (7)$$

$$= \text{tr} \left[\exp(\ln \rho) \exp((\alpha - 1) \ln(\rho_A \otimes \mathbf{1})) \right] \quad (8)$$

$$\geq \text{tr} \left[\exp(\ln(\rho) + (\alpha - 1) \ln(\rho_A \otimes \mathbf{1})) \right] \quad (9)$$

$$\geq \text{tr} \left[\exp(\alpha \ln \rho) \right] = \text{tr}(\rho^\alpha), \quad (10)$$

where Eq.(9)-(10) is implied by the reduction criterion (2) since the logarithm is operator monotone [21] and the exponential function is monotone under the trace. The latter can be seen by noting that for any A hermitian, $P \geq 0$ and $B = (A + \epsilon P)$ with $\epsilon \geq 0$:

$$\frac{\partial}{\partial \epsilon} \text{tr}(e^B) = \text{tr}(e^B P) \geq 0. \quad (11)$$

Hence $\text{tr}(e^B) \geq \text{tr}(e^A)$ is implied by $B \geq A$.

For $0 \leq \alpha < 1$ the reduction criterion can immediately be applied since $f(A) = A^r$ is an operator decreasing function for $-1 \leq r \leq 0$, $A \geq 0$ (cf. [22]) and thus

$$\text{tr}(\rho_A^\alpha) = \text{tr}[\rho(\rho_A^{\alpha-1} \otimes \mathbf{1})] \leq \text{tr}(\rho^\alpha), \quad (12)$$

which completes the proof. $\square$

For negative values of α , $T_\alpha(\rho) \geq 0$ holds for any state no matter if it is entangled or not. This can be shown by noting that for $\{|a\rangle\}$ being an eigenbasis of ρ_A

$$\text{tr}(\rho_A^\alpha) = \sum_a \langle a | \rho_A | a \rangle^\alpha = \sum_a \left[\sum_i \langle a \otimes i | \rho | a \otimes i \rangle \right]^\alpha \quad (13)$$

$$\leq \sum_{a,i} \langle a \otimes i | \rho | a \otimes i \rangle^\alpha \leq \text{tr}(\rho^\alpha), \quad (14)$$

where Eq.(13-14) uses that $(\sum_i b_i)^\alpha \leq \sum_i b_i^\alpha$ holds for $b_i \geq 0$, $\alpha \leq 0$, and the last inequality is implied by the convexity of negative powers.

We note that the von Neumann case $S_1(\rho_A) \leq S_1(\rho)$ resp. $T_1(\rho) \geq 0$ also follows from the reduction criterion due to the operator monotonicity of the logarithm. Hence, separability criteria based on conditional Tsallis-Rényi entropies for all positive values of the entropic parameter are implied by the reduction criterion and therefore certainly weaker than the latter. In particular they cannot be sufficient.

This raises the question about the resolution that separability criteria, which are based on the spectrum of a state and its reductions, can have in general. The next two sections are devoted to the somewhat disillusioning answer.

III. SEPARABLE COUNTERPARTS FOR WERNER STATES

Werner states [4] have always played an important and paradigmatic role in quantum information theory. Their characteristic property is that they commute with all unitaries of the form $U \otimes U$ and they can be expressed as

$$\rho(p) = (1-p)\frac{P_+}{r_+} + p\frac{P_-}{r_-}, \quad 0 \leq p \leq 1, \quad (15)$$

where P_+ (P_-) is the projector onto the symmetric (antisymmetric) subspace of $\mathbb{C}^d \otimes \mathbb{C}^d$ and $r_{\pm} = \text{tr}[P_{\pm}] = \frac{d^2 \pm d}{2}$ are the respective dimensions. Werner showed that these states are entangled iff $p > \frac{1}{2}$ independent of the dimension d . The following proposition shows however, that none of these entangled states for odd dimension d can be detected by any separability criterion, which is based on the spectrum of the state and its reductions.

Proposition 2 *Any entangled state in $\mathbb{C}^d \otimes \mathbb{C}^d$ with maximal chaotic reductions and eigenvalues having multiplicities which are multiples of d , has a separable isospectral counterpart, which is locally undistinguishable as it has the same reductions.*

Proof: Let us consider a special basis of maximally entangled states in $\mathbb{C}^d \otimes \mathbb{C}^d$ [23]:

$$|\Psi_{jk}\rangle = \frac{1}{\sqrt{d}} \sum_{n=1}^d \exp\left(\frac{2\pi i}{d} jn\right) |n, n \oplus k\rangle, \quad (16)$$

where $j, k = 1, \dots, d$ and $\oplus$ means addition modulo d . Any equal weight combination of all states of the form (16), which belong to the same value of k , is then a projector onto a separable state since

$$\begin{aligned} P_k &= \sum_{j=1}^d |\Psi_{jk}\rangle \langle \Psi_{jk}| \\ &= \frac{1}{d} \sum_{j,n,m=1}^d \exp\left[\frac{2\pi i}{d} j(n-m)\right] |n, n \oplus k\rangle \langle m, m \oplus k| \\ &= \sum_{n=1}^d |n\rangle \langle n| \otimes |n \oplus k\rangle \langle n \oplus k| \end{aligned} \quad (17)$$

is an equal weight combination of product states. Here we have used that $\frac{1}{d} \sum_{j=1}^d \exp\left[\frac{2\pi i}{d} j(n-m)\right] = \delta_{n,m}$. Moreover, the reductions of the respective states P_k/d are maximally chaotic, i.e. $\rho_A = \mathbf{1}/d$, just as the reductions of any maximally entangled state.

If we now have a state with multiplicities being multiples of d we can replace the projectors onto its eigenspaces with sufficiently many projectors of the form P_k . The resulting state will then be again a convex combination of product states, i.e., separable, having the same spectrum and maximal chaotic reductions. $\square$

For the case of Werner states we note that the unitary invariance of the state $\rho(p)$ in Eq. (15) implies that its reductions are $\rho_A = \mathbf{1}/d$. Moreover $\rho(p)$ has two eigenvalues $(1-p)/r_+$ and p/r_- with multiplicities r_+ , r_- which are indeed multiples of d in odd dimensions.

Following Proposition 2 we can now construct a state

$$\rho'(p) = \frac{(1-p)}{r_+} \sum_{k=1}^{r_+/d} P_k + \frac{p}{r_-} \sum_{l=1}^{r_-/d} P_{l+r_+/d}, \quad (18)$$

which has then both, the same spectrum and the same reductions as $\rho(p)$. However, as convex combination of separable states it is itself separable for any $0 \leq p \leq 1$.

IV. THE TWO QUBIT CASE

For the case of two qubits the antisymmetric subspace is one dimensional and the Werner state $\rho(p)$ is therefore entangled iff it has an eigenvalue larger than one half. It was essentially shown by Bennett et al. [24] that this is indeed sufficient to decide separability for two qubit states with maximal chaotic reductions, such that these states cannot have separable counterparts.

However, the following will show that separability is in general not encoded in the spectrum of a two qubit state and its reductions.

It is often advantageous to express two qubit states in terms of the matrix

$$R_{ij} = \text{tr}(\rho \sigma_i \otimes \sigma_j), \quad i, j = 0, \dots, 3 \quad (19)$$

with $\sigma_0 = \mathbf{1}$ and $\sigma_{1,2,3}$ being the Pauli matrices. Normalization then requires $R_{0,0} = 1$ and the reductions are determined by the vectors $R_{0,i}$ and $R_{i,0}$. However, positivity of the respective state has to be verified separately.

Let us now consider a controlled phase gate operation described by the unitary

$$U = |0\rangle\langle 0| \otimes \mathbf{1} + |1\rangle\langle 1| \otimes \sigma_3 \quad (20)$$

$$= \frac{1}{2}(\mathbf{1} + \sigma_3) \otimes \mathbf{1} + \frac{1}{2}(\mathbf{1} - \sigma_3) \otimes \sigma_3 \quad (21)$$

which applies a phase operation on the target bit if the source bit is in the state 1 and leaves it unchanged if the source bit is in the state 0. Controlled phase gates as well as controlled NOT gates can be considered as universal basic building blocks for future quantum computers since together with single qubit gates they can be used to implement an arbitrary quantum circuit (cf. [25]).

The operation U interchanges the matrix elements $R_{1,0} \leftrightarrow R_{1,3}$, $R_{0,1} \leftrightarrow R_{3,1}$ and $R_{1,1} \leftrightarrow R_{2,2}$ since

$$U(\sigma_1 \otimes \mathbf{1})U^* = (\sigma_1 \otimes \sigma_3), \quad (22)$$

$$U(\mathbf{1} \otimes \sigma_1)U^* = (\sigma_3 \otimes \sigma_1), \quad (23)$$

$$U(\sigma_1 \otimes \sigma_1)U^* = (\sigma_2 \otimes \sigma_2). \quad (24)$$

If we now choose the state such that $R_{0,1} = R_{1,0} = R_{3,1} = R_{1,3} = r$ take on the same values, $R_{1,1} = \frac{1}{2}$ and the others vanish, then the controlled phase gate will leave the reductions unchanged and simply interchange $R_{1,1} \leftrightarrow R_{2,2}$:

$$R = \begin{pmatrix} 1 & r & 0 & 0 \\ r & \frac{1}{2} & 0 & r \\ 0 & 0 & 0 & 0 \\ 0 & r & 0 & 0 \end{pmatrix} \xrightarrow{U} R' = \begin{pmatrix} 1 & r & 0 & 0 \\ r & 0 & 0 & r \\ 0 & 0 & \frac{1}{2} & 0 \\ 0 & r & 0 & 0 \end{pmatrix} \quad (25)$$

These matrices correspond to positive states for $|r| \leq \frac{3}{8}$. Moreover, since the partial transposition only changes the signs of the σ_2 components, the initial state is equal to its partial transpose and therefore separable on the whole interval. This argumentation fails however for the state $\rho' = U\rho U^*$, which corresponds to R' , and in fact the determinant of its partial transpose is negative and the state is thus entangled for $\frac{\sqrt{3}}{8} < |r| \leq \frac{3}{8}$.

Hence, the controlled phase gate operation can produce an entangled state out of a separable one, which has the same reductions and due to the unitarity of the operation the same spectrum.

It is worth mentioning that although ρ' is entangled for $\frac{\sqrt{3}}{8} < |r| \leq \frac{3}{8}$ it does not violate the CHSH Bell inequality [26]. It can be shown that $T_2 \geq 0$ is indeed stronger than the Bell inequality for two qubit systems [12]. Hence, a similar example with one state being separable and the other violating the CHSH inequality cannot exist. This is however no longer true for higher dimensional systems.

V. CONCLUSION

Deciding whether a given quantum state is entangled or not is still one of the big open problems in the theory of quantum information. We discussed to what extent criteria based on the spectrum of a state and its reductions can decide separability. We showed that criteria, which are based on entropic inequalities using conditional Tsallis and Rényi entropies, are definitely weaker than the reduction criterion but nevertheless valid for any value of the entropic parameter. Moreover, the fact that there exists a separable counterpart for any entangled Werner state in odd dimensions shows that there is only a rather loose relation between the spectra of a state and its reductions on the one hand and its entanglement properties on the other.

Since the latter must not depend on the choice of local bases, it is sufficient to look at local unitary invariants of a state in order to decide its separability. Surprisingly, the most efficient separability criterion is still based on just one rather simple invariant: the smallest eigenvalue

of the partial transpose. One remaining question in this context would therefore be: how can other (easy calculable) invariants provide information about the separability of a state, which is not yet encoded in the smallest eigenvalue of its partial transpose?

ACKNOWLEDGEMENT

The authors would like to thank R.F. Werner for useful discussions. Funding by the European Union project EQUIP (contract IST-1999-11053) and financial support from the DFG (Bonn) is gratefully acknowledged.

-
- [1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [2] E. Schrödinger, Naturwissenschaften **23**, 807-812; 823-828; 844-849 (1935).
 - [3] J.S. Bell, Physics **1**, 195 (1964).
 - [4] R.F. Werner, Phys. Rev. A **40**, 4277 (1989).
 - [5] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer, 1995).
 - [6] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).
 - [7] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 1 (1996).
 - [8] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **80**, 5239 (1998).
 - [9] M. Horodecki and P. Horodecki, Phys. Rev. A **59**, 4206 (1999).
 - [10] Note that throughout this paper we omit equations corresponding to the reduction with respect to the second subsystem $\rho_B = \text{tr}_A(\rho)$. However, all the equations hold in a symmetric form for both reductions and the presented ones should be read as representatives for either of the two.
 - [11] N. Cerf and C. Adami, Phys. Rev. Lett. **79**, 5194 (1997).
 - [12] R. Horodecki, P. Horodecki, and M. Horodecki, Phys. Lett. A **210**, 377 (1996); R. Horodecki and M. Horodecki, Phys. Rev. A **54**, 1838 (1996).
 - [13] S. Abe and A.K. Rajagopal, Physica A **289**, 157 (2001).
 - [14] S. Abe and A.K. Rajagopal, Phys. Rev. A **60**, 3461 (1999).
 - [15] C. Tsallis, S. Lloyd, and M. Baranger, Phys. Rev. A **63**, 042104 (2001).
 - [16] A. Vidiella-Barranco, Phys. Lett. A, 260, 335 (1999).
 - [17] A.K. Rajagopal and R.W. Rendell, quant-ph/0106050 (2001).
 - [18] P. Horodecki, J.A. Smolin, B.M. Terhal, and A.V. Thapliyal, quant-ph/9910122 (1999); B.M. Terhal, quant-ph/0101032 (2001).
 - [19] Abe and Rajagopal already tried to prove this proposition for separable states in [13]. However, they failed since they wrongly supposed that all the states in the decomposition in Eq.(1) mutually commute.

- [20] M. Ohya and D. Petz, *Quantum Entropy and Its Use* (Springer 1993).
- [21] K. Löwner, Math. Z. **38**, 177 (1934).
- [22] A.W. Marshall and I. Olkin, *Inequalities: Theory of Majorization and Its Applications* (Academic Press 1979).
- [23] R.F. Werner, quant-ph/0003070 (2000).
- [24] C.H. Bennett, D.P. DiVincenzo, J.A. Smolin, W.K. Wootters, Phys. Rev. A **54**, 3824 (1996).
- [25] D.P. DiVincenzo, Proc. R. Soc. London A **454**, 261 (1998).
- [26] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).